# **Lesson 1.2**: Problem Solving (Part 2) 3. Language of Real Mathematics Problem Mathematical **Problem** Check Check 5. Solution **Problem Solving Model**

# 5 Categories of Problems

- Direct Translation problems where we must translate from English into Mathematics by using key words in the verbal description.
- Mixture problems where two or more quantities are combined in some fashion.
- Geometry problems where the unknown quantities are related through geometric formulas.
- Uniform Motion problems where an object travels at a constant speed.
- Work Problems problems where two or more entities join forces to complete a job.

# Steps for Solving Problems with Mathematical Models

- Step 1: Identify what you are looking for.
- Step 2: Give Names to the Unknowns.
- Step 3: Translate the Problem into the Language of Mathematics.
- Step 4: Solve the Equation(s) Found in Step 3.
- Step 5: Check the Reasonableness of your Answer.
- Step 6: Answer the Question (in a complete sentence).

Lesson 1.2: Problem Solving (Part 2) Step 2:

Ex 1: I= Prt; where I = Interest (simple interest),
P = Principal, r = rate of interest, and
t = time (in years)

Supposed that Kenzie has a credit card balance of \$2800. Each month, the credit card company charges 14% annual simple interest on any oustanding balances. What is the interest that Kenzie will be charged on this loan after one month? What is Kenzie's credit card balance after one month?

\$1273: \$\sum\_{==}^{2800} (.44)(\frac{1}{12})\$\$

P= 
$$\frac{2.800}{12}$$
 Step 4:  $I = \frac{32.67}{12}$ 

t=  $\frac{1}{12}$  Step 6: The interest is \$\frac{1}{12}\$ Step 6: they balance will be \$\frac{1}{12}\$ 2832.67.

Ex 2: Suppose that Ethan has a car loan of \$6500. The bank charges 6% annual simple interest. What is the interest charge on Ethan's car loan after 1 month?

$$I = ?$$
 Step 2:  $I = INterest$   
 $P = 6500$  Step 3:  $I = 6500(.06)(12)$   
 $r = .06$  Step 4:  $I = 325$ 

t=(定) Step 6: The interest after I month

# Mixing Problems

Mixture problems are problems in which two or more items are combined to form a third item. There are a number of different kinds of mixture problems, but they all follow the basic idea.

Portion from item A + Portion from item B = Whole or Total

**Ex 3:** Mitch has \$15,000 to invest. His goal is to obtain an overall annual rate of return of 9% or \$1350 annually. His financial advisor recommends that he invest some of the money in corporate bonds that pay 12% and the rest in government-backed Treasury bonds paying 4%. How much should be placed in each investment in order for Mitch to achieve his goal?

# Step 1: Identify

Mitch needs to figure out how much to put in corporate bonds vs. Treasury bonds in order to get \$1350 in interest.

# Step 2: Name

Let's have b represent the amount Mitch invests in corporate bonds. How can we represent the amount Mitch invests in Treasury bonds? b+t=15000

**Ex 3:** Remember: Mitch has \$15,000 to invest. His goal is to obtain an overall annual rate of return of 9% or \$1350 annually. Corporate bonds that pay 12% and Treasury

bonds pay 4%.

# Step 3: Translate

So now, we need a mathematical model.

	Principal (\$)	Rate (%)	Time (Yr)	Interest (\$)
Corporate Bond	ь	0.12	1	0.12b
Treasury Bond	15,000 – b	0.04	1	0.04(15,000 – b)
Total	15,000	0.09	1	0.09(15000) = \$1350

Interest from Corporate bonds + Interest from Treasury bonds = \$1350

**Ex 3:** Remember: Mitch has \$15,000 to invest. His goal is to obtain an overall annual rate of return of 9% or \$1350 annually. Corporate bonds that pay 12% and Treasury bonds pay 4%.

Step 4: Solve
$$.12b + .04(15000 - 6) = 1350$$

$$.12b + 600 - .04b = 1350$$

$$.08b + 600 - 600 + 1350$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

$$-600 - 600 + 15000 - 9345$$

**Ex 3:** Remember: Mitch has \$15,000 to invest. His goal is to obtain an overall annual rate of return of 9% or \$1350 annually. Corporate bonds that pay 12% and Treasury bonds pay 4%.

Step 5: Check

#### Step 6: Answer the Question

Mitch invested 97375 in corp. bonds & \$5625 in treasury bonds.

**Ex 4:** The manager of a coffee shop wishes to form a new blend of coffee. She wants to mix Sumarta Beans, known for their strong, distinctive taste, that sell for \$12/lb with milder Brazilian Beans that sell for \$8/lb to get 50 lbs of the new blend. The new blend will sell for \$9/lb and there will be NO difference in revenue from selling the new blend vs. selling the beans separately. How many lbs. of the Sumatra and Brazilian beans are required?

# Step 1: Identify

This is a mixture problem becuase we want to know how many pounds of each (Sumatra and Brazilian) beans we need to make a new blend.

# Step 2: Name

Let's have **s** represent the amount of Sumatra Beans that are required. So what will the # of lbs be for the Brazilian beans?

Ex 4:

Step 3: Translate

	Price (\$/lb)	Number of lbs.	Revenue
Sumatra	12	s	12s
Brazilian	8	50-s	8(50-s)
Blend	9	50	9(50) = \$450

So now, we need a mathematical model.

Revenue from Sumatra + Revenue from Brazilian = Revenue of New Blend

$$12s + 8(so-3) = 450$$

# Ex 4:

Step 4: Solve
$$|2S + 8(50-S) = 450$$

$$|2S + 400 - 8S = 450$$

$$|2S + 400 - 8S = 450$$

$$|2S + 400 - 450$$

# Step 5: Check

# Step 6: Answer the Question

She heeds 12.51bs of sumatrabeans and 37.51bs of the Brazilian Beans.

# <u>Uniform Motion Problems</u>

Objects that move at a constant velocity are said to be in uniform motion. When the average velocity of an object is known, it can be interpreted as its constant velocity.

For example, a car traveling at an average velocity of 40 miles/hour is in uniform motion.

If and object moves at an average speed r, the distance d covered in time t is given by:

$$d = rt$$

**Ex 5:** Austin and Dillon decide to have a 10-mile race. Austin can run at an average speed of 12 miles/hour while Dillon can run at an average speed of 10 miles/hour. To "even things up", Austin agrees to give Dillon a head start of 0.15 hour. When will Austin catch up to Dillon?

Step 1: Identify

when will they meet?

Step 2: Name

t=time

**Ex 5:** Austin and Dillon decide to have a 10-mile race. Austin can run at an average speed of 12 miles/hour while Dillon can run at an average speed of 10 miles/hour. To "even things up", Austin agrees to give Dillon a head start of 0.15 hour. When will Austin catch up to Dillon?

#### Step 3: Translate

	Rate (mph)	Time (hrs)	Distance (Miles)	
Austin	12	H	12t	
Dillon	10	Ftis	10 (t+.15)	
12t = 10(t+.15)				



#### Ex 5:

# Step 3: Translate


So now, we need a mathematical model.

	Rate (mph)	Time (hrs)	Distance (Miles)
Austin	12	t	12t
Dillon	10	t + 0.15	10(t + 0.15)

Distance Austin runs = Distance Dillon runs

Ex 5:

Step 4 : Solve



Step 5: Check

Step 6: Answer the Question

# Homework:

```
Pg. 76:
30, 37, 40, 42, 43, 49-51 all,
53, 57, 59, 61, 63, 65, 68
```

(14 problems)