

Lesson 47: Solving Log Equations

By the end of the lesson, we will be able to:

- ~ Solve log equations
- ~ Solve equations with more than one log



Lesson 47: Solving Log Equations

One way to solve log equations is to rewrite in exponential form. We use this method when there is log expression on one side of the equation, and a number on the other side.

Converting between exponential and log form:

$$\text{If } \log_b x = y, \text{ then } b^y = x$$

Lesson 47: Solving Log Equations

Examples: Convert to exponent form and solve.

a.) $\log_6 x = 2$

b.) $\log_4(3x) = 3$

Lesson 47: Solving Log Equations

Examples: Convert to exponent form and solve.

$$c.) 5 + 2 \ln x = 9$$

Lesson 47: Solving Log Equations

If there is more than one log expression in the equation, the logs must first be condensed.

Example: $\log_2 x + 4 \log_2 3 = 4$

Lesson 47: Solving Log Equations

Important!!!

Always Check Your Solutions!

Some solutions may be extraneous.

A solution is extraneous if...

Lesson 47: Solving Log Equations

A solution is extraneous if...

it is a solution of the simplified form of an equation that does not satisfy the original equation.

inside of log cannot be zero or negative

Lesson 47: Solving Log Equations

Examples: Condense and solve.

a.) $\log(x + 2) + \log(x - 1) = 1$

Lesson 47: Solving Log Equations

Examples: Condense and solve.

$$b.) \log_5(4x + 7) - \log_5 x = 2$$

Lesson 47: Solving Log Equations

Another way to solve log equations is to use the following property of equality:

Property of equality for log functions:

$$\text{If } \log_b x = \log_b y, \text{ then } x = y$$

If there is a single log on both sides, with the same base, then the arguments can be set equal.

Lesson 47: Solving Log Equations

Examples:

a.) $\log_3(3x - 5) = \log_3(x + 7)$

Lesson 47: Solving Log Equations

Examples:

$$b.) \log(x^2 - 6) = \log x$$

Lesson 47: Solving Log Equations

If there is more than one log expression in the equation, the logs must first be condensed.

Examples:

a.) $\log_2(x + 3) - \log_2(2x - 1) = \log_2 2$

Lesson 48: Exponential Growth and Decay

Real life situations involving exponential growth or decay can be modeled using the equation:

Exponential Growth and Decay:

$$y = Pe^{rt}$$

where y is final amount, P is initial amount (Principal), r is the growth rate, and t is time.

Lesson 48: Exponential Growth and Decay

This equation $y = Pe^{rt}$ is used when we are compounding interest continually (always).

But sometimes, we only compound interest annually (once a year). We use this equation when we do that: $y = P(1 + r)^t$

Lesson 48: Exponential Growth and Decay

In this equation $y = P(1 + r)^t$, r stands for rate.

~ If r is positive, we are growing.

~ If r is negative, we are decaying
(decreasing).

Lesson 48: Exponential Growth and Decay

$$y = P(1 + r)^t$$

Examples: Give the percent growth or decay represented by the following equation.

B. $y = 3000(1.65)^t$

Lesson 48: Exponential Growth and Decay

$$y = P(1 + r)^t$$

Examples: Give the percent growth or decay represented by the following equation.

c. $y = 4000(.75)^t$

Lesson 48: Exponential Growth and Decay

$$y = P(1 + r)^t$$

Examples: Give the percent growth or decay represented by the following equation.

D. $y = 3500 \left(\frac{3}{4}\right)^t$

Lesson 48: Exponential Growth and Decay

Another application of exponential equations is **radioactive decay**. Radioactive elements decay over time, and the rate of decay is constant. Instead of a decay rate (r), the decay is often given as the **half life** of the substance. Half life is the amount of time it takes for half of the mass to decay.

When given the half life of an element, the decay rate (r) can be found using the formula:

Decay rate when given half life:

$$r = \frac{-\ln 2}{\text{half life}}$$

Lesson 48: Exponential Growth and Decay

$$y = Pe^{rt}$$

Examples:

Element X has a half-life of 2000 years. A sample starts with 830 grams.

H. How many grams will be left in 105 years?

$$y =$$

$$P =$$

$$r =$$

$$t =$$

$$r = \frac{-\ln 2}{\text{half life}}$$

Lesson 48: Exponential Growth and Decay

$$y = Pe^{rt}$$

Examples:

Element X has a half-life of 2000 years. A sample starts with 830 grams.

I. How long until 200 grams remain?

$$y =$$

$$P =$$


$$r =$$

$$t =$$

$$r = \frac{-\ln 2}{\text{half life}}$$

When are logs used?

The bel (symbol B) is a unit of measure which is the base -10 logarithm of ratios, such as power levels and voltage levels. It is mostly used in telecommunication, electronics, and acoustics. The Bel is named after telecommunications pioneer Alexander Graham Bell. The decibel (dB), equal to 0.1 bel, is more commonly used.



When are logs used?

The Richter scale measures earthquake intensity on a base -10 logarithmic scale.





When are logs used?

In astronomy, the apparent magnitude measures the brightness of stars logarithmically, since the eye responds approximately logarithmically to brightness.



When are logs used?

Musical intervals are measured logarithmically as semitones (half steps up or down).



Lesson 47: Solving Log Equations

By the end of the lesson, we will be able to:

- ~ Solve log equations
- ~ Solve equations with more than one log



Lesson 48: Exponential Growth and Decay

By the end of the lesson, we will be able to:

- ~ Solve application problems with exponentials and logs
 - * Exponential Growth problems
 - * Exponential Decay problems

Homework:

Journal 47 + Journal 48
(Skip #5-10 - you already
did them.)

Assignment 48